

CASE REPORT

L. D. R. Thompson

Langerhans cell histiocytosis isolated to the thyroid gland

Received: 19 April 1995

Abstract Isolated Langerhans cell histiocytosis of the thyroid is an extremely rare occurrence, with only one case previously reported. A case of Langerhans cell histiocytosis isolated to the thyroid gland associated with lymphocytic thyroiditis is presented and clinical implications are discussed.

Key words Langerhans cell histiocytosis · Thyroid gland · Histiocytosis X · Differential diagnosis

Introduction

Langerhans cell histiocytosis (LCH) is the appellation now applied to a rare spectrum of disorders that includes eosinophilic granuloma, Letterer-Siwe disease and Hand-Schuller-Christian disease [7, 18]. LCH is a disorder of histiocytes of uncertain etiology and generally affects bone, lung, lymph nodes and central nervous system. Adults are infrequently affected. Although there has been involvement of the thyroid gland reported in the literature as part of systemic disease [1–6, 8–17], only a report by Coode and Shaikh [2] has described isolated thyroid disease. The patient herein reported presented with localized thyroid disease. This case report is part of an ongoing review of thyroid Langerhans cell histiocytosis.

Case report

A 32-year-old Caucasian male presented to an endocrinologist with a 3–4 week history of a thyroid mass. His thyroxine, tri-

iodothyronine uptake and thyroid-stimulating hormone level were all within normal limits. A thyroid scan was performed, and demonstrated a 2-cm cold nodule in the right lobe of the thyroid gland. The patient declined fine needle aspiration for cytologic evaluation, and a surgical hemi-thyroidectomy was performed as an excisional biopsy and definitive therapy.

The excised thyroid tissue was red-brown, homogeneous in consistency and contained a small focal nodular mass occupying its central portion. Histologic examination of hematoxylin and eosin stained preparations demonstrated a small focal collection of histiocytes with a small number of eosinophils (Fig. 1). The lobular, indented and folded nuclei were surrounded by a microvacuolated clear to eosinophilic cytoplasm (Fig. 2). The histiocytic infiltrate had destroyed thyroid follicles, effacing the normal glandular architecture (Fig. 3). A mild lymphocytic thyroiditis was present in the remainder of the gland. Immunohistochemical stains were then performed, with S-100 protein and lysozyme reacting positively. Since the disease focus was small, additional immunohistochemical reactions and electron microscopy could not be performed. However, all findings were consistent with LCH.

The patient has since remained free of systemic LCH for the past 4 years, and has not had any other thyroid abnormalities. Thyroid function tests have remained normal to date.

Discussion

To date, isolated reports about patients with LCH thyroid involvement have included patients with concurrent systemic disease [1–6, 8–17]. Additionally, all of the reported cases seem to have an associated lymphocytic thyroiditis in addition to hypothyroidism and diabetes insipidus. The latter may imply that there is involvement of the pituitary gland, but has often not been so stated in the reports published.

The occurrence of hypothyroidism in LCH could have a number of causes: it may be related to pituitary involvement, result from destruction of thyroid follicles by histiocytic infiltrate, and/or be due to lymphocytic thyroiditis. It is unknown which factor(s) is/are actually occurring.

Diagnosis of LCH is predicated on tissue histopathology, since usual thyroid function studies may be normal and nuclear scans show only cold nodules (Fig. 4). Although the histologic findings are typical for LCH as it occurs elsewhere in the body (as in our case), the lobulated and grooved or indented nuclei can cause some con-

L. D. R. Thompson
Department of Otolaryngic and Endocrine Pathology,
Armed Forces Institute of Pathology, Building 54, Room 2025,
6825 16th Street, NW, Washington, DC 20306–6000, USA

The opinions or assertions contained herein are the private views of the author and are not to be construed as official or as reflecting the views of the Department of the Navy or the Department of Defense




Fig.1 Histopathology of thyroid gland demonstrating lymphocytic thyroiditis and focal Langerhans cell histiocytosis. Hematoxylin and eosin stain, $\times 30$




Fig.3 Histopathology of thyroid gland specimen with typical follicular epithelial effacement, Langerhans cell histiocytes and eosinophils. Hematoxylin and eosin stain, $\times 400$




Fig.2 Histopathology of thyroid gland specimen showing lobulated, indented and folded nuclei surrounded by pale to eosinophilic cytoplasm. A few eosinophils are seen. Hematoxylin and eosin stain, $\times 900$




Fig.4 Radionuclide scan with radioactive ^{131}I labelling demonstrating a cold nodule of the right lobe of the thyroid gland that was proven histologically to be Langerhans cell histiocytosis

fusion with papillary carcinoma. The typical infiltrate may also be obscured by the lymphocytic infiltration that occurs with lymphocytic thyroiditis. Careful histologic examination as well as confirmatory testing for S-100 protein and lysozyme or another macrophage-associated marker by immunohistochemical staining [such as KP-1 (CD68)] will lead to an accurate diagnosis.

The usual treatment for systemic disease associated with LCH often involves radiation or chemotherapy, while it would seem from the reported case in the literature and our case that localized disease is best managed successfully through surgery alone. It is therefore important to ascertain whether the thyroid involvement by LCH is part of systemic disease; if it is localized, curative surgical intervention may be all that a patient with localized thyroid disease requires.

References

1. Avery ME, McAfee JG, Guild HG (1957) The course and prognosis of reticuloendotheliosis. *Am J Med* 22: 636–652
2. Coode PE, Shaikh MU (1988) Histiocytosis X of the thyroid masquerading as thyroid carcinoma. *Hum Pathol* 19: 239–241
3. Deméocq F, Boespflug O, Vanlieferinghen P, Jourdan JE, Lesec G, Malpuech G (1982) Localisation thyroïdienne d'une histiocytose X. *Arch Fr Pédiatr* 39: 449–453
4. Goldstein N, Layfield L (1991) Thyromegaly secondary to simultaneous papillary carcinoma and histiocytosis X. *Acta Cytol* 35: 422–426
5. Lahey ME, Rallison ML, Hilding DA, Ater J (1986) Involvement of the thyroid in histiocytosis X. *Am J Pediatr Hematol Oncol* 8: 257–259
6. Latorre H, Kenney FM, Lahey ME, Drash A (1974) Short stature and growth hormone deficiency in histiocytosis X. *J Pediatr* 85: 813–816
7. Lichtenstein L (1953) Histiocytosis X: integration of eosinophilic granuloma of bone, Letterer-Siwe disease and Schüller-Christian disease as related manifestations of a single nosologic entity. *Arch Pathol* 56: 84–102
8. Ma JT, Wang C, Ho FC, Lam KS, Yeung RT (1985) Primary hypothyroidism and essential hyponatremia in a patient with histiocytosis X. *Aust NZJ Med* 15: 72–74
9. Mahzoon S, Wood MG (1980) Multifocal eosinophilic granuloma with skin ulceration. *Arch Dermatol* 116: 218–220
10. Panayotopoulos E, Concouris L, Roussis S, Sinacos Z (1964) Granulome éosinophile de la glande thyroïde avec diabète insipide. *Presse Med* 72: 153–154
11. Rothman JG, Synder PJ, Utiger RD (1978) Hypothalamic endocrinopathy in Hand-Schüller-Christian disease. *Ann Intern Med* 88: 512–513
12. Schafer EL (1949) Nonlipid reticulo-endotheliosis: Letterer-Siwe's disease. *Am J Pathol* 25: 49–73
13. Sinisi AA, Criscuolo T, Palombini L, Bellastella A, Faggiano M (1986) Thyroid localization in adult histiocytosis X. *J Endocrinol Invest* 9: 417–420
14. Teja K, Sabio H, Langdon DR, Johanson AJ (1981) Involvement of the thyroid gland in histiocytosis X. *Hum Pathol* 12: 1137–1139
15. Turiaf J, Battesti J, Basset F, Racadot J, Poirier J, Hugues F (1971) Histiocytose X de l'adulte à localisations pulmonaires, thyroïdiennes, méningo-encéphaliques et hypophysothalamiques. *Semin Hop Paris* 47: 2923–2932
16. Whitaker C, Kenny FM, Fetterman GH (1971) Asymptomatic goiter as an expression of a rare disease. *Clin Pediatr* 10: 290–292
17. Wilkins L (1965) The diagnosis and treatment of endocrine disorders in childhood and adolescence, 3rd edn. Thomas, Springfield, Ill, p 157
18. Writing group of the Histiocyte Society (1987) Histiocytosis syndromes in children. *Lancet* I: 208–209